

University of Wollongong

Research Online

Faculty of Informatics - Papers (Archive)

Faculty of Engineering and Information
Sciences

1973

Kronecker products and BIBDS

Jennifer Seberry

University of Wollongong, jennie@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/infopapers>



Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Seberry, Jennifer: Kronecker products and BIBDS 1973.
<https://ro.uow.edu.au/infopapers/946>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Kronecker products and BIBDS

Abstract

Recursive constructions are given which permit, under conditions described in the paper, a (v, b, r, k, λ) -configuration to be used to obtain a (v', b', r', k, λ) -configuration.

Although there are many equivalent definitions we will mean by a (v, b, r, k, λ) -configuration or BIBD that $(0, 1)$ -matrix A of size $v \times b$ with row sum r and column sum k satisfying

$$AA^T = (r - \lambda)I + \lambda J$$

where, as throughout the remainder of this paper, I is the identity matrix and J the matrix with every element $+1$ whose sizes should be determined from the context or by a subscript (J_n is square of order n).

Disciplines

Physical Sciences and Mathematics

Publication Details

Jennifer Seberry Wallis, Kronecker products and BIBDS, Journal of Combinatorial Theory, Ser. A., 14, (1973), 248-252.

Note

Kronecker Products and BIBDs

JENNIFER WALLIS

Faculty of Mathematics, University of Newcastle, N.S.W. 2308, Australia

Communicated by M. Hall, Jr.

Received April 13, 1971

Recursive constructions are given which permit, under conditions described in the paper, a (v, b, r, k, λ) -configuration to be used to obtain a (v', b', r', k, λ) -configuration.

Although there are many equivalent definitions we will mean by a (v, b, r, k, λ) -configuration or *BIBD* that $(0, 1)$ -matrix A of size $v \times b$ with row sum r and column sum k satisfying

$$AA^T = (r - \lambda)I + \lambda J$$

where, as throughout the remainder of this paper, I is the identity matrix and J the matrix with every element $+1$ whose sizes should be determined from the context or by a subscript (J_n is square of order n).

In the case of block matrices, $(X)_{ij}$ and (X_{ij}) mean the matrix whose (i, j) -th block is X ; for example, $(T^{i-j})_{ij}$ is the matrix whose (i, j) -th block is T^{i-j} . We define the *Kronecker product* of two matrices $A = (a_{ij})$ of order $m \times n$ and B of any order as the $m \times n$ block matrix

$$A \times B = (a_{ij}B)_{ij}.$$

For more details the reader is referred to Marshall Hall [1].

We will use T for the circulant matrix of order q given by

$$T = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 1 & 0 & 0 & \cdots & 0 & 0 & 1 \end{bmatrix}.$$

For q a prime we have shown in Jennifer Wallis [3] that

$$Q = (T^{(i-1)(j-1)})_{ij}$$

satisfies

$$\begin{aligned} QQ^T &= qI_q \times I_q + (J_q - I_q) \times J_q, \\ J \cdot Q &= qJ. \end{aligned}$$

We are concerned with the existence of a $(0, 1)$ matrix Q of size $mv \times v^2$ which satisfies

$$\begin{aligned} QQ^T &= vI_m \times I_v + (J_m - I_m) \times J_v, \\ JQ &= mJ; \end{aligned} \quad (1)$$

if such a matrix exists we will say $\mathbf{P}(m, v)$ holds. Thus the result cited above shows that $\mathbf{P}(q, q)$ holds for any prime q ; we also showed in [3] that $\mathbf{P}(q, q)$ holds for any prime power q . Further, it was proved in [4] that $\mathbf{P}(m, v)$ holds if and only if there exists a set of $m - 2$ mutually orthogonal Latin squares of order v , and that a $(0, 1)$ matrix Q satisfying (1) must have the form

$$Q = \begin{bmatrix} E \\ A \\ \vdots \\ A_{m-1} \end{bmatrix},$$

where E and the A_i are of size $v \times v^2$ and have constant row sums v and column sums 1. From the latter fact it is clear that if Q satisfies (1) then the matrix formed by deleting A_n and subsequent blocks satisfies (1) with m replaced by n , so

$$\mathbf{P}(m, v) \Rightarrow \mathbf{P}(n, v) \quad \text{when } n < m.$$

If we are referring to $\mathbf{P}(m, v)$, then Q , E and A_i will mean the matrices just mentioned.

MAIN THEOREM

We shall exploit the following theorem, which is a generalization of Lemma 6 of [3]:

THEOREM 1. *Suppose B is a (v, b, r, k, λ) -configuration and suppose R is a $(0, 1)$ matrix of size $lv \times tv^2$ satisfying*

$$\begin{aligned} RR^T &= a_1vI_l \times I_v + a_2(J_l - I_l) \times J_v, \\ JR &= kJ, \end{aligned} \quad (2)$$

where a_2 divides λ . Then necessarily $la_1 = kt$ and $(l-1)a_2 = (k-1)a_1$, and

$$[I_l \times B \mid R, R, \dots, R] \quad (\lambda/a_2 \text{ copies of } R)$$

is an $(lv, lb + \lambda tv^2/a_2, r + \lambda a_1 v/a_2, k, \lambda)$ -configuration.

Proof. By summing the entries of R in two ways we obtain $la_1 = kt$. It is easy to check that the matrix exhibited is the required configuration; one of the standard necessary conditions for a (v, b, r, k, λ) -configuration is $\lambda(v-1) = r(k-1)$; substituting in the parameters of the configuration we constructed we have

$$(l-1)a_2 = (k-1)a_1.$$

In the particular case where $t = a_1 = a_2 = 1$ and $k = l$, the existence of a suitable R satisfying (2) is simply $\mathbf{P}(k, v)$.

COROLLARY 2. *If there exist a (v, b, r, k, λ) -configuration and a set of $k-2$ mutually orthogonal Latin squares of order v , then there is a $(kv, kb + \lambda v^2, r + \lambda v, k, \lambda)$ -configuration.*

EXAMPLE. Suppose $v = p_1^{a_1} p_2^{a_2} \cdots p_n^{a_n}$ is a decomposition of v into powers of distinct primes, and suppose

$$k \leq \min_i (p_i^{a_i}) + 1.$$

Then there is a set of $k-2$ mutually orthogonal Latin squares of order v [1, p. 192], so the existence of a (v, b, r, k, λ) -configuration for this k and v implies the existence of a $(kv, kb + \lambda v^2, r + \lambda v, k, \lambda)$ -configuration.

EXAMPLE. Hanani [2] has shown (in terms of Latin squares) that $\mathbf{P}(5, v)$ always holds when $v \geq 52$ and $\mathbf{P}(7, v)$ always holds when $v \geq 63$, so Corollary 2 can be applied in the corresponding cases.

FIRST APPLICATION

Suppose q is a prime and ω is a primitive q -th root of unity. T is of order q . Define a $q \times q$ matrix P by

$$P = (p_{ij}), \quad p_{ij} = \omega^{(i-1)(j-1)}.$$

Now define square matrices S_{ij} , $i = 1, 2, \dots, q^s$ and $j = 1, 2, \dots, q^s$ where

s is any positive integer, as follows: if the (i, j) element of the Kronecker product of s copies of P is ω^a , then $S_{ij} = T^a$.

Assume that $\mathbf{P}(k, v)$ holds for some $k \leq q^s$. Write

$$R = \begin{bmatrix} S_{11} \times E & S_{12} \times E & \cdots & S_{1q^s} \times E \\ S_{21} \times A_1 & S_{22} \times A_1 & \cdots & S_{2q^s} \times A_1 \\ \vdots & \vdots & \ddots & \vdots \\ S_{k1} \times A_{k-1} & S_{k2} \times A_{k-1} & \cdots & S_{kq^s} \times A_{k-1} \end{bmatrix}.$$

R is a $(0, 1)$ matrix of size $kvq \times v^2q^{s+1}$, and it is readily shown that

$$RR^T = q^s v I_k \times I_{qv} + q^{s-1} (J_k - I_k) \times J_{qv},$$

so R satisfies (2) with v replaced by vq , $l = k$ and $t = a_1 = a_2 = q^{s-1}$. So we have proved the following:

THEOREM 3. *Suppose there exists a (qv, b, r, k, λ) -configuration, where q is a prime, and suppose $\mathbf{P}(k, v)$ holds. If s is a positive integer such that q^{s-1} divides λ and $k \leq q^s$, then there exists a $(kvq, kb + \lambda q^2 v^2, r + \lambda qv, k, \lambda)$ -configuration.*

Corollaries can easily be constructed using the examples in the preceding section.

SECOND APPLICATION

Suppose q is a prime and suppose $\mathbf{P}(k, v)$ holds where $k \leq q + 1$. The matrices I and T will be of order q .

We consider the $(0, 1)$ block matrix P ,

$$P = \begin{bmatrix} I \times A_1 & I \times A_1 & I \times A_1 & \cdots & I \times A_1 \\ I \times A_2 & T \times A_2 & T^2 \times A_2 & \cdots & T^{q-1} \times A_2 \\ I \times A_3 & T^2 \times A_3 & T^4 \times A_3 & \cdots & T^{2(q-1)} \times A_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ I \times A_{k-1} & T^{k-2} \times A_{k-1} & T^{2(k-2)} \times A_{k-1} & \cdots & T^{(k-2)(q-1)} \times A_{k-1} \end{bmatrix},$$

which is a $(k-1) \times q$ array of $qv \times qv^2$ blocks. Write E' for $[I \times E, I \times E, \dots, I \times E]$, there being q copies of $I \times E$, and denote by $T^i \cdot P$ the result of multiplying the first of the two components of every block entry of P by T^i . Then

$$R = \begin{bmatrix} E' & E' & E' & \cdots & E' \\ P & T \cdot P & T^2 \cdot P & \cdots & T^{q-1} \cdot P \end{bmatrix}$$

is a $(0, 1)$ matrix of suitable size which satisfies (2) with $l = k$, v replaced by qv and $a_1 = a_2 = t = q$. Hence we have

THEOREM 4. *If $P(k, v)$ holds and there is a (qv, b, r, k, λ) -configuration, where q is a prime not less than $k - 1$ and q divides λ , then there exists a $(kqv, kb + \lambda q^2 v^2, r + \lambda qv, k, \lambda)$ -configuration.*

Again corollaries can be formed at will.

REFERENCES

1. M. HALL, JR., "Combinatorial Theory," Blaisdell, Waltham, Mass., 1967.
2. H. HANANI, On the number of orthogonal Latin squares, *J. Combinatorial Theory* **8** (1970), 247-271.
3. J. WALLIS, Some results on configurations, *J. Austral. Math. Soc.* **12** (1971), 378-384.
4. W. D. WALLIS, A remark on Latin squares and block designs, *J. Austral. Math. Soc.* **13** (1972), 205-207.